

## **Exploiting T<sub>1</sub>-dispersion using human-scale fast field-cycling MRI**

David Lurie, Lionel Broche, Gareth Davies, Mary-Joan MacLeod, P. James Ross

*School of Medicine, Medical Sciences & Nutrition, University of Aberdeen, Scotland, UK*

Much of the contrast in conventional MRI arises from disease-induced changes in T<sub>1</sub>. Extra information could be obtained from T<sub>1</sub>-dispersion measurements (T<sub>1</sub> versus magnetic field), but this information is invisible to standard MRI scanners, since they operate only at fixed magnetic field (e.g. 1.5 T, 3.0 T). We have developed Fast Field-Cycling Magnetic Resonance Imaging (FFC-MRI) to exploit T<sub>1</sub>-dispersion as a potential biomarker of disease, with the aim of increasing diagnostic potential.

T<sub>1</sub>-dispersion is typically measured using FFC, by switching the magnetic field rapidly between levels during the pulse sequence; relaxation occurs at the (low) evolution field while detection is always at the same (higher) detection field. Thus, a single instrument can be used to measure T<sub>1</sub> over a wide range of magnetic field strengths. FFC-MRI obtains spatially-resolved T<sub>1</sub>-dispersion data, by collecting images at a range of evolution fields.

We have built a variety of FFC-MRI equipment, including two whole-body human scanners, operating at detection fields of 0.06 T and 0.2 T. Recent work has focused on speeding up FFC-MRI using rapid pulse sequences, as well as the investigation of methods to measure T<sub>1</sub>-dispersion at ultra-low magnetic fields.

In vitro measurements in our laboratory have shown that FFC can detect changes in human cartilage induced by osteoarthritis; we have also demonstrated that T<sub>1</sub>-dispersion is sensitive to cancer-induced changes in breast tissues. We are exploring clinical applications and have imaged patients who have had an ischaemic stroke; the affected brain tissues are seen as hyper-intense regions in ultra-low-field (200 μT) FFC images.

This presentation will cover the main techniques used in FFC-MRI and will summarise current and potential bio-medical applications.

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 668119 (project "IDentIFY").